

## Spring Wheat Facts (NASS-ID)

- Harvested Area
  - 2020 – 495,000 acres
- Average Yield
  - 2020 – 91 bu/A
- Production
  - 2020 – 45,045,000 bu
    - 60 lb = 1 bu

## Growth and Development

Using Feekes Growth Scale - Vegetative stage is through Feekes 5, reproductive stage begins at 6

- Germination –when seed is exposed to adequate moisture, oxygen, and temp.
- Seedling growth – until 9 or more leaves have unfolded
- Tillering – from 1 to 5 tillers
- Stem elongation – starting from detection of 1<sup>st</sup> node
- Booting – flag leaf sheath extended to first visible awns
- Inflorescence emergence – Spikelet visible to complete emergence
- Anthesis – 5 to 7 days after heading, beginning to completion of flowering
- Milk – kernel development to late milk
- Dough – early (mealy), soft to hard dough
- Ripening – kernel approaches harvest moisture (hard dough to harvest ready)

## Rotation and Seeding

- Wheat grows well in rotation – not recommended after corn or small grains when alternatives are available
- Good seed-to-soil contact is needed
- Seed depth should be 1 to 1.5 in under irrigation and good soil moisture
- Row-spacing of 6 to 8 in with commercial drills provides uniform distribution of seed
- Seeding rate\* depends on seed size
  - Irrigated: 1 – 1.2 million seeds per acre (65 to 120 lb/A)
  - Dryland: 700,000 seeds per acre (55 to 90 lb/A)

\*Increased seeding rates recommended with delayed planting or poor seed bed.
- Optimum germination - when soil temperature is between 55 and 75°F

## Optimum Planting Date Estimates

| <u>Location</u>         | <u>Timing</u>            |
|-------------------------|--------------------------|
| Treasure Valley         | Late Feb to mid-March    |
| Magic Valley            | Mid-March to early April |
| Upper Snake River Plain | Late March to late April |

## Irrigation

- Time to meet evapotranspiration (ET) and seasonal crop needs

- Greatest yield reduction occurs with moisture stress at:
  - Tillering
  - Boot to flowering
- Evapotranspiration (ET)
  - ~ 15 to 19 in of water
  - Peak ET occurs in mid-June to mid-July at heading and decreases after soft dough
- Water Holding Capacity (WHC) – the amount of water held in soil for crops
  - Soil texture WHC estimates
    - Loamy > 2 in/ft
    - Sandy loams 1 to 2 in/ft
    - Sandy < 1 in/ft
- Available Soil Moisture (ASM) – the difference between existing soil moisture content and permanent wilting point
  - ASM can be estimated by subtracting ET from the WHC if the soil profile WHC and soil moisture lost to ET are known
- Center Pivot Systems
  - Early season - supply soil root zone with moisture
  - Late season, pivot may not supply sufficient water to keep up with ET in which case **additional soil water reserves will be needed**

# 2020 S. Idaho Soft White Spring Wheat Quick Facts

- Surface Irrigation Systems
  - 1<sup>st</sup> irrigation should occur at 50% ASM (earlier on sandy soil)
  - At least 50% ASM maintained from tillering to soft dough

## Fertilization

- Soil Sampling
  - One to two weeks prior to planting
  - 0- to 12-in and 12- to 24-in sample depth for nitrogen (N) and sulfur (S) separated by depth
  - 0- to 12-in for other nutrients
- Estimate of Nitrogen rate – 2.0-2.5 units N/bu yield based on:
  - Inorganic soil test N
  - Mineralizable N from OM = 30-60 lbs N/A (estimated typically at 45 lb N/A)
  - Crop residues
    - Potato/sugarbeet/onion residue is accounted for by soil test
    - Alfalfa provides 40 to 80 lb N/A beyond soil testing
    - Small grain residue – ADD 15 lb N for each ton of residue returned to the soil (up to 50 lb N/A)
  - Application timing
    - Loamy soil – single preplant or 40% preplant, 60% at tillering
    - Sandy soil – split 40% preplant, 60% at tillering

- No additional N recommended after tillering for SWS
- Phosphorus (P, P<sub>2</sub>O<sub>5</sub>) - Pounds of P<sub>2</sub>O<sub>5</sub> applied based on soil test and percent free lime.

| <u>Olsen Soil Test (0-12 in)</u> | <u>Percent free lime</u>                |     |     |     |
|----------------------------------|---|-----|-----|-----|
|                                  | 0                                       | 5   | 10  | 15  |
| ppm                              | lbs P <sub>2</sub> O <sub>5</sub> /acre |     |     |     |
| 0                                | 240                                     | 280 | 320 | 360 |
| 5                                | 160                                     | 200 | 240 | 280 |
| 10                               | 80                                      | 120 | 160 | 200 |
| 15                               | 0                                       | 40  | 80  | 120 |
| 20                               | 0                                       | 0   | 0   | 40  |

- Potassium (K, K<sub>2</sub>O)
  - Response can be expected in soil with <75 ppm K (0-12 in sample)
- Sulfur (S, SO<sub>4</sub>)
  - 0- to 24 in sample depth
  - At < 10 ppm S (or <35 lb/A) and low-S irrigation water
    - 20 to 40 lbs/A of sulfate-based fertilizer can result in yield response
- Other important nutrients: Chloride (Cl), Fe, Mn, Fe, Zn, Cu, B

## **Growth Regulators**

Ethephon (Cerone) and/or Palisade  
Apply at labeled rates and timing to reduce lodging, plant height

## **Common Diseases**

Stripe rust, Fusarium head blight (FHB), root rots (Fusarium crown rot, Rhizoctonia, take-all), cereal cyst nematode, bacterial blight, loose smut, seedling blight (Pythium) and other nematodes

## **Common Insect Pests**

Aphids, cereal leaf beetle, thrips, Haanthen barley mealybug, wireworms, armyworms, cutworms

## **Common Weeds**

- Annuals: wild oat, green foxtail, kochia, common lambsquarters, redroot pigweed, feral rye, wild buckwheat, and various mustards
- Perennials: Canada thistle, field bindweed, quackgrass

**Prepared by: J.M. Marshall, C.W.**

**Rogers, A. Rashed, O. Walsh, X. Liang,**

**O.S. Walsh, and A. Adjesiwor**

## **References:**

- Brown, B. Walsh, O. 2016. Planting Dates in Wheat Production in Southern Idaho. University of Idaho AES BUL 906.
- Hagerty and Smiley. 2017. A Field Guide for Diagnosing Common Wheat Maladies of the PNW. PNW 698.
- Brown, B., Stark, J., Westermann, D. 2001. Southern Idaho Fertilizer Guide: Irrigated Spring Wheat. University of Idaho AES BUL 828.